

# CONTENTS

Opinions: Open Science		<p><b>Prehabilitation Programs Before Radical Prostatectomy: An Interesting Approach that Merits Further Evaluation</b> <span style="float: right;">12</span></p> <p><i>H. Isbarn</i></p>
Brief Correspondence		<p><b>Salvage Radiotherapy Following Partial Gland Ablation for Prostate Cancer: Functional and Oncological Outcomes</b> <span style="float: right;">1</span></p> <p><i>Y. Qaoud, P. Bettoli, N. Sanmamed-Salgado, J.O. Herrera-Caceres, M.B. Berjaoui, K. Lajkosz, H. Goldberg, D.T.S. Woon, Z. Glase, S. Ghai, A. Finelli, P. Chung, N. Perlis, N. Fleshner, A. Berlin</i></p> <p><b>Confirmation by Early Oncologic Outcomes After Surgery of the Accuracy of Intermediate-risk Prostate Cancer Classification Based on Magnetic Resonance Imaging Staging and Targeted Biopsy</b> <span style="float: right;">5</span></p> <p><i>C. Manceau, J.-B. Beauval, M. Lesourd, C. Almeras, J.-R. Gautier, M. Soulié, G. Loison, A. Salin, C. Tollon, B. Malavaud, M. Roumiguié, G. Ploussard</i></p> <p><b>What Stone-formers Should Know About Vitamin C and D Supplementation in the COVID-19 Era</b> <span style="float: right;">9</span></p> <p><i>J.A. Khusid, W.M. Atallah, N. Kyprianou, M. Gupta</i></p> <p><b>One-day Prehabilitation Program Before Robotic Radical Prostatectomy in Daily Practice: Routine Feasibility and Benefits for Patients and Hospitals</b> <span style="float: right;">14</span></p> <p><i>G. Ploussard, G. Loison, C. Almeras, J.-R. Gautier, P. Cazali, C. Tollon, J.-B. Beauval, A. Salin</i></p> <p><b>Overexpression of Placental Growth Factor in Stromal Cells from Benign Prostatic Hyperplasia: Another Piece in the Puzzle?</b> <span style="float: right;">29</span></p> <p><i>C.M. Devlin, F.M. Frame, L.K. Archer, V.M. Mann, H. Walker, M.S. Simms, D.M. Berney, A.T. Collins, N.J. Maitland</i></p> <p><b>Can Exercise Adaptations Be Maintained in Men with Prostate Cancer Following Supervised Programmes? Implications to the COVID-19 Landscape of Urology and Clinical Exercise</b> <span style="float: right;">47</span></p> <p><i>P. Lopez, D.R. Taaffe, R.U. Newton, N. Spry, T. Shannon, M. Frydenberg, F. Saad, D.A. Galvão</i></p>
Original Articles		<p><b>Identification and Validation of Leucine-rich <math>\alpha</math>-2-glycoprotein 1 as a Noninvasive Biomarker for Improved Precision in Prostate Cancer Risk Stratification</b> <span style="float: right;">51</span></p> <p><i>I.J. Guldvik, V. Zuber, P.R. Braadland, H.H. Grytli, H. Ramberg, W. Lilleby, B. Thiede, M. Zucknick, F. Saatcioglu, R. Gislefoss, R. Kvåle, A. George, H. Grönberg, F. Wiklund, D.E. Neal, V.J. Gnanapragasam, K.A. Taskén, I.G. Mills</i></p> <p>High blood levels of leucine-rich <math>\alpha</math>-2-glycoprotein 1 (LRG1) is unfavourable in patients with newly diagnosed high-risk or metastatic prostate cancer. Implementation of a blood test for LRG1 alongside standard risk stratification schemes could provide higher precision in treatment decisions.</p>

	<p><b>Impact of Health-related Quality of Life and Prediagnosis Risk of Major Depressive Disorder on Treatment Choice in Low- and Intermediate-Risk Prostate Cancer</b> 69</p> <p><i>H. Riskin-Jones, T. Grogan, A. Kishan, A. Raldow</i></p> <p>Depressive symptoms and physical health-related quality of life are associated with the type of treatment prostate cancer patients receive, emphasizing the importance of considering these factors when counseling patients and developing decision aids.</p>
Urothelial Cancer	<p><b>Fibroblast Growth Factor Receptor 3 Mutation as a Prognostic Indicator in Patients with Urothelial Carcinoma: A Systematic Review and Meta-analysis</b> 61</p> <p><i>S. Khalid, B.M. Basulaiman, J. Emack, C.M. Booth, I. Duran, A.G. Robinson, D. Berman, M. Smoragiewicz, E. Amir, F.E. Vera-Badillo</i></p> <p>Fibroblast growth factor receptor 3 (FGFR3) mutation is not associated with a worse survival outcome in urothelial carcinoma. This is relevant as clinical trials are being conducted to assess the response of FGFR inhibitors in urothelial carcinoma.</p>
Kidney Cancer	<p><b>Impact of Positive Surgical Margins After Partial Nephrectomy</b> 41</p> <p><i>J.A.M. Carvalho, P. Nunes, E. Tavares-da-Silva, B. Parada, R. Jarimba, P. Moreira, E. Retroz, R. Caetano, V. Sousa, A. Cipriano, A. Figueiredo</i></p> <p>Positive surgical margins after partial nephrectomy were associated with a high risk of pathological lesions and lower surgeon volume. There was no impact on survival.</p>
Reconstructive Urology	<p><b>Robotic Management of Pelvic Lipomatosis—Experience with Difficulties Encountered and the Techniques to Successful Outcomes</b> 33</p> <p><i>J. SanjayPrakash, T. Mathisekaran, N. Jain, S. Bafna, R. Paul, N. Selvaraj</i></p> <ul style="list-style-type: none"> <li>•Robotic management of pelvic lipomatosis causing bilateral hydroureteronephrosis has good surgical outcome.</li> <li>•Imaging cannot always reliably detect this condition.</li> <li>•The choice of surgical technique purely depends on the surgeon's comfort, leaving logistic issues aside.</li> </ul>
Education	<p><b>Impact of COVID-19 on Clinical and Academic Urological Practice: A Survey from European Association of Urology Section of Uro-technology</b> 22</p> <p><i>A. Heinze, P. Umari, M. Basulto-Martínez, R. Suárez-Ibarrola, E. Liatsikos, J. Rassweiler, S. Guven, A.S. Gözen</i></p> <p>As situations similar to the coronavirus disease 2019 (COVID-19) pandemic may recur, we must be prepared to react quickly and effectively. It is important that urology departments look after the well-being of their patients, while protecting their staff and continuing with academic/research activities.</p>
Case Report	<p><b>Percutaneous Biopsy Tract Seeding in a Patient with Muscle-invasive Bladder Cancer</b> 17</p> <p><i>A. Gusev, S. Greenberg, S. Dave, A. Sobieh, J. Yates</i></p>