

CONTENTS

<p>Opinion: Open Science</p> 	<p>Treatment Intensification for Low-risk Metastatic Hormone-sensitive Prostate Cancer: The Time Is Now</p> <p><i>R.R. McKay</i></p>	<p>41</p>
<p>Brief Correspondence</p> 	<p>Diagnosis of Clinically Significant Prostate Cancer Diagnosis Without Histological Proof in the Prostate-specific Membrane Antigen Era: The Jury Is Still Out</p> <p><i>M. Wenzel, B. Hoeh, P. Mandel, F.K. Chun</i></p>	<p>50</p>
<p>Open Horizon</p> 	<p>Intravesical Ty21a Treatment of Non-muscle-invasive Bladder Cancer Shows a Good Safety Profile</p> <p><i>I. Lucca, L. Derré, V. Cesson, P. Bohner, F. Crettenand, S. Rodrigues-Dias, F. Dartiguenave, A. Masnada, C. Teixeira-Pereira, S. Benmerzoug, M. Chevalier, S. Domingos-Pereira, S. Nguyen, L. Polak, A. Schneider, B. Roth, P. Jichlinski, D. Nardelli-Haeffliger</i></p>	<p>55</p>
<p>Reviews</p> 	<p>Hormonal Intensification Should Start at the Low-risk Stage in Metastatic Prostate Cancer</p> <p><i>S.B. Jazayeri, L.F. Cooley, A. Srivastava, N. Shore</i></p>	<p>38</p>
<p>Original Articles</p> 	<p>Role of Pediatric Ureteral Access Sheath and Outcomes Related to Flexible Ureteroscopy and Laser Stone Fragmentation: A Systematic Review of Literature</p> <p><i>F. Ripa, T. Tokas, S. Griffin, S. Ferretti, A. Bujons Tur, B.K. Somani</i></p> <p>Ureteral access sheath (UAS) placement during flexible ureteroscopy and laser lithotripsy in the pediatric population is safe and associated with a low rate of ureteric injuries. Further studies are needed to clarify its indications and to assess the clinical outcomes of UAS placement.</p>	<p>90</p>
<p>Prostate Cancer</p>	<p>Impact of Ureteral Stent Material on Stent-related Symptoms: A Systematic Review of the Literature</p> <p><i>M. Boeykens, E.X. Keller, A. Bosio, O.J. Wiseman, P. Contreras, E. Ventimiglia, M. Talso, A. Pietropaolo, T. Tailly, V. De Coninck</i></p> <p>Silicone DJ stents, and by extension soft DJ stents, appear to reduce stent-related symptoms. No definitive conclusion can be drawn because of the lack of quality evidence. A standard for measuring physical stent properties should be developed.</p>	<p>108</p>
<p>Prostate Cancer</p>	<p>Real-world Treatment Sequencing in Patients with Metastatic Castration-resistant Prostate Cancer: Results from the Prospective, International, Observational Prostate Cancer Registry</p> <p><i>A. Bjartell, L. Costa, G. Kramer, B. Zurawski, L. Galli, P. Werbrouck, T. Ecke, O. Parikh, M. Bennamoun, C. Garcia Freire, A. Peer, B. Ljungberg, I. Cicin, E. Smith, M. Lukac, R. Wapenaar, S. Chowdhury</i></p> <p>Patient characteristics appear to influence the first-line treatment of metastatic castration-resistant prostate cancer. Patients older and with lower Gleason scores tended to receive abiraterone, those younger and with more advanced disease tended to receive docetaxel, and those with fewer metastases and more favourable performance tended to receive enzalutamide. The benefit to patients from these observations remains unknown.</p>	<p>12</p>

	<p>Reliability of Ancestry-specific Prostate Cancer Genetic Risk Score in Four Racial and Ethnic Populations 23</p> <p>Z. Shi, J. Zhan, J. Wei, S. Ladson-Gary, C.-H. Wang, P.J. Hulick, S. L. Zheng, K.A. Cooney, W.B. Isaacs, B.T. Helfand, B.L. Koelsch, J. Xu</p> <p>Based on 1 013 652 individuals from four ancestry groups (European, Hispanic, African, and East Asian), we demonstrated the reliability of the ancestry-specific genetic risk score for prostate cancer (concordance with the observed risk) and suitability for genetic testing.</p>
	<p>Estetrol Prevents Hot Flushes and Improves Quality of Life in Patients with Advanced Prostate Cancer Treated with Androgen Deprivation Therapy: The PCombi Study 59</p> <p>Y. Zimmerman, M. Frydenberg, H. van Poppel, R.J.A. van Moorselaar, E.P.M. Roos, D.M. Somford, T.A. Roeleveld, T.D. de Haan, H.H.E. van Melick, Y. Reisman, J. Krijgh, F.M.J. Debruyne, H.J.T. Coelingh Bennink</p> <p>Androgen deprivation therapy (ADT) for advanced prostate cancer is accompanied by side effects affecting healthrelated quality of life (HRQL). Cotreatment with ADT and high-dose estetrol almost completely prevented the occurrence of hot flushes and improved HRQL and well being in the PCombi trial.</p>
	<p>Low Blood Levels of LRG1 Before Radical Prostatectomy Identify Patients with High Risk of Progression to Castration-resistant Prostate Cancer 68</p> <p>I.J. Guldvik, P.R. Braadland, S. Sivanesan, H. Ramberg, G. Kristensen, P. Tennstedt, A. Røder, T. Schlomm, V. Berge, L.M. Eri, W. Lilleby, I.G. Mills, K.A. Taskén</p> <p>Measurement of LRG1 in blood before radical prostatectomy may provide guidance on follow-up regimens and treatment decisions.</p>
Kidney Cancer	<p>Impact of Partial Nephrectomy and Percutaneous Cryoablation on Short-term Health-related Quality of Life—A Prospective Comparative Cohort Study 99</p> <p>T. Junker, L. Duus, B.S.B. Rasmussen, N. Azawi, L. Lund, B. Nørgaard, O. Graumann</p> <p>Patients receiving percutaneous cryoablation (PCA) had lower baseline scores on physical functioning but were comparable with those receiving partial nephrectomy (PN) with regard to Global Health Status.</p> <p>Significant differences in health-related quality of life were reported from baseline to 14 d postoperatively between PN and PCA, but were no longer evident after 90 d.</p>
Sexual Medicine	<p>Sexual Activity and Dyspareunia After Pelvic Organ Prolapse Surgery: A 5-Year Nationwide Follow-up Study 81</p> <p>O. Wiheraari, P. Karjalainen, A.-M. Tolppanen, N. Mattsson, K. Nieminen, J. Jalkanen</p> <p>This 5-yr follow-up study demonstrates that pelvic organ prolapse (POP) surgery improves sexual activity and reduces dyspareunia. Multiple factors, such as preoperative pain, prior POP surgery, and prolapse stage, may be associated with postoperative dyspareunia.</p>
Education	<p>Evidence-based Medicine: Perceptions, Attitudes, and Skills Among European Urology Residents 44</p> <p>D.A. González Padilla, J.D. Subiela, D.M. Carrion, F. Esperto, J. Gómez Rivas, S. Khadhour, G. Mantica, A. Mattigk, B. Pradere, M. Rodríguez Socarrás, E. Segui Moya, J.Y.-C. Teoh, L. Tortolero-Blanco, T. Uçar, P. Dahm</p> <p>Formal training in evidence-based medicine is lacking in most urology residency programs. Perceptions of and attitudes to evidence-based medicine are very positive among European urology residents.</p>

Surgery in Motion: Open Science	"Light green up": Indocyanine Green Fluorescence Imaging-guided Robotic Bilateral Inguinal Lymphadenectomy by the Hypogastric Subcutaneous Approach for Penile Cancer	1
	<i>P. Yuan, K. Yao, Z. Zhou, J. Liu, C. Li, W. Hou, Y. Tang, S. Hu, L. Wang</i>	
	Indocyanine green fluorescence imaging-guided robotic bilateral inguinal lymphadenectomy via a hypogastric subcutaneous approach is safe and effective for selected patients with penile cancer. This technique may be beneficial for increasing the number of dissected inguinal lymph nodes.	
Nightmares in Urology: Open Science	Technical Tips in Managing Large Median Lobes During Robot-assisted Radical Prostatectomy	32
	<i>V.G. Wagaskar, O. Zaytoun, P. Kale, A. Pedraza, D. Busby, A. Reddy, A. Tewari</i>	
	Robot-assisted radical prostatectomy in patients with a large median lobe is a challenging procedure with a steep learning curve. It is associated with increased operative time and estimated blood loss, but with similar oncological and continence outcomes. Understanding the key steps to facilitate bladder neck dissection is vital to avoid serious intraoperative events and to maximize outcomes.	
Trial Protocol	Clinical Trial Protocol for PERFECT: A Randomised Controlled Trial Comparing the Efficiency and Tolerance of Transperineal Fusion Versus Transrectal Imaging-targeted Prostate Biopsies (CCAFU-PR1 Study)	76
	<i>A. Touzani, G. Fiard, E. Barret, R. Renard-Penna, A. Salin, B. Pradère, F. Rozet, J.-B. Beauval, B. Malavaud, G. Giannarini, P. Colin, M. Rouprêt, G. Ploussard</i>	
	This study evaluates the efficiency of the transrectal and transperineal approaches for prostate biopsy in terms of detecting significant cancer, and compares the safety between these two approaches.	
Case Report	Low-energy Shockwave Therapy in the Management of Wound Healing Following Fournier's Gangrene	8
	<i>J.J. Rassweiler, W. Scheitlin, A.S. Goezen, M.-C. Rassweiler-Seyfried</i>	
Letters to the Editor	Reply to Julian Chavarriaga and Robert Hamilton's Letter to the Editor re: Manolis Pratsinis, Christian Fankhauser, Katerina Pratsinis, et al. Metastatic Potential of Small Testicular Germ Cell Tumors: Implications for Surveillance of Small Testicular Masses. Eur Urol Open Sci 2022;40:16–8. Should We Be Afraid of Surveillance? Clinically Meaningful Reasons Why Offering Surveillance for Incidentally Detected Small Testicular Masses Remains a Safe Approach	52
	<i>M. Pratsinis, C. Rothermundt, C. Fankhauser</i>	
	Re: Manolis Pratsinis, Christian Fankhauser, Katerina Pratsinis, et al. Metastatic Potential of Small Testicular Germ Cell Tumors: Implications for Surveillance of Small Testicular Masses. Eur Urol Open Sci 2022;40:16–18.	53
	<i>J. Chavarriaga and R. Hamilton</i>	
Corrigendum	Corrigendum re "21 – Urine exosomal miRNA profiles implicate the key role of TP53 in bladder cancer development" [Eur Urol Open Sci 2022;40(Suppl 1):S31–2]	31
	<i>E. Bitina-Barlote, J. Plonis, J.K. Vasionoks, E. Vjaters, J. Gardovskis, Z. Daneberga, E. Miklaševičs, M. Nakazawa-Miklaševiča</i>	